

Percentage of Adults 65 Years and Older Vaccinated Against Influenza in 2003*

State	Yes	No
Alabama	70.2%	29.8%
Alaska	66.5%	33.5%
Arizona	68.9%	31.1%
Arkansas	71.0%	29.0%
California	72.5%	27.5%
Colorado	74.2%	25.8%
Connecticut	74.3%	25.7%
Delaware	70.0%	30.0%
District of Columbia	63.0%	37.0%
Florida	65.9%	34.1%
Georgia	67.0%	33.0%
Hawaii	76.4%	23.6%
Idaho	70.3%	29.7%
Illinois	63.3%	36.7%
Indiana	66.1%	33.9%
Iowa	77.5%	22.5%
Kansas	70.8%	29.2%
Kentucky	69.1%	30.9%
Louisiana	68.3%	31.7%
Maine	74.8%	25.2%
Maryland	68.4%	31.6%
Massachusetts	74.9%	25.1%
Michigan	67.5%	32.5%
Minnesota	80.3%	19.7%
Mississippi	69.0%	31.0%
Missouri	69.9%	30.1%

State	Yes	No
Montana	72.8%	27.2%
Nebraska	73.6%	26.4%
Nevada	60.0%	40.0%
New Hampshire	73.9%	26.1%
New Jersey	67.2%	32.8%
New Mexico	72.4%	27.6%
New York	68.0%	32.0%
North Carolina	68.8%	31.2%
North Dakota	73.0%	27.0%
Ohio	68.0%	32.0%
Oklahoma	75.8%	24.2%
Oregon	70.5%	29.5%
Pennsylvania	69.1%	30.9%
Rhode Island	76.2%	23.8%
South Carolina	69.3%	30.7%
South Dakota	77.9%	22.1%
Tennessee	69.1%	30.9%
Texas	67.7%	32.3%
Utah	74.8%	25.2%
Vermont	74.1%	25.9%
Virginia	69.6%	30.4%
Washington	73.4%	26.6%
West Virginia	69.1%	30.9%
Wisconsin	72.1%	27.9%
Wyoming	72.6%	27.4%

***Healthy People 2010 Goal = 90%**

Source: Division of Adult and Community Health, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, [Behavioral Risk Factor Surveillance System Online Prevalence Data](#), 1995-2003.