

Percentage of Adults 65 Years and Older Who Have Ever Received a Pneumococcal Vaccination*

State	Yes	No
Alabama	61.4%	38.6%
Alaska	59.6%	40.4%
Arizona	65.5%	34.5%
Arkansas	61.8%	38.2%
California	65.2%	34.8%
Colorado	69.1%	30.9%
Connecticut	64.5%	35.5%
Delaware	67.4%	32.6%
District of Columbia	50.1%	49.9%
Florida	64.4%	35.6%
Georgia	60.5%	39.5%
Hawaii	69.4%	30.6%
Idaho	67.2%	32.8%
Illinois	56.6%	43.4%
Indiana	61.5%	38.5%
Iowa	71.4%	28.6%
Kansas	60.3%	39.7%
Kentucky	59.6%	40.4%
Louisiana	64.2%	35.8%
Maine	64.8%	35.2%
Maryland	62.0%	38.0%
Massachusetts	69.4%	30.6%
Michigan	62.7%	37.3%
Minnesota	73.0%	27.0%
Mississippi	61.7%	38.3%
Missouri	61.1%	38.9%

State	Yes	No
Montana	69.1%	30.9%
Nebraska	64.8%	35.2%
Nevada	63.2%	36.8%
New Hampshire	69.3%	30.7%
New Jersey	62.4%	37.6%
New Mexico	63.9%	36.1%
New York	61.7%	38.3%
North Carolina	66.6%	33.4%
North Dakota	71.2%	28.8%
Ohio	64.7%	35.3%
Oklahoma	68.6%	31.4%
Oregon	71.7%	28.3%
Pennsylvania	66.1%	33.9%
Rhode Island	69.3%	30.7%
South Carolina	63.0%	37.0%
South Dakota	63.7%	36.3%
Tennessee	60.8%	39.2%
Texas	62.0%	38.0%
Utah	66.2%	33.8%
Vermont	66.1%	33.9%
Virginia	65.2%	34.8%
Washington	68.6%	31.4%
West Virginia	63.8%	36.2%
Wisconsin	66.7%	33.3%
Wyoming	70.4%	29.6%

***Healthy People 2010 Goal = 90%**

Source: Division of Adult and Community Health, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, [Behavioral Risk Factor Surveillance System Online Prevalence Data](#), 1995-2003.