



## Percentage of Adults 65 Years and Older Vaccinated Against Influenza in 2005\*

State	Yes	No
<b>Nationwide</b>	<b>65.5%</b>	<b>35.5%</b>
Alabama	60.8%	39.2%
Alaska	61.1%	38.9%
Arizona	62.5%	37.5%
Arkansas	65.2%	34.8%
California	65.9%	34.1%
Colorado	74.2%	25.8%
Connecticut	71.1%	28.9%
Delaware	65.8%	34.2%
District of Columbia	54.7%	45.3%
Florida	55.6%	44.4%
Georgia	60.8%	39.2%
Hawaii	72.1%	27.9%
Idaho	63.9%	36.1%
Illinois	55.9%	44.1%
Indiana	64.0%	36.0%
Iowa	71.7%	28.3%
Kansas	66.0%	34.0%
Kentucky	62.4%	37.6%
Louisiana	62.4%	37.6%
Maine	67.8%	32.2%
Maryland	59.3%	40.7%
Massachusetts	69.8%	30.2%
Michigan	67.2%	32.8%
Minnesota	78.2%	21.8%
Mississippi	61.5%	38.5%

State	Yes	No
Missouri	61.7%	38.3%
Montana	69.5%	30.5%
Nebraska	72.6%	27.4%
Nevada	53.0%	47.0%
New Hampshire	70.2%	29.8%
New Jersey	63.4%	36.6%
New Mexico	68.0%	32.0%
New York	61.8%	38.2%
North Carolina	65.5%	34.5%
North Dakota	70.1%	29.9%
Ohio	64.7%	35.3%
Oklahoma	73.2%	26.8%
Oregon	68.9%	31.1%
Pennsylvania	59.3%	40.7%
Rhode Island	67.2%	32.8%
South Carolina	60.9%	39.1%
South Dakota	76.3%	23.7%
Tennessee	61.6%	38.4%
Texas	61.6%	38.4%
Utah	69.6%	30.4%
Vermont	66.3%	33.7%
Virginia	66.8%	33.2%
Washington	67.8%	32.2%
West Virginia	63.6%	36.4%
Wisconsin	71.8%	28.2%
Wyoming	72.9%	27.1%

**\*Healthy People 2010 Goal = 90%**

Source: Division of Adult and Community Health, National Center for Chronic Disease Prevention and Health Promotion, Centers for Disease Control and Prevention, Behavioral Risk Factor Surveillance System Online Prevalence Data, 1995-2005.